

Science

 AAAS

The Hallucinogen N,N-Dimethyltryptamine (DMT) Is an Endogenous Sigma-1 Receptor Regulator

Dominique Fontanilla, *et al.*

Science **323**, 934 (2009);

DOI: 10.1126/science.1166127

The following resources related to this article are available online at www.sciencemag.org (this information is current as of February 12, 2009):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/323/5916/934>

Supporting Online Material can be found at:

<http://www.sciencemag.org/cgi/content/full/323/5916/934/DC1>

This article **cites 29 articles**, 11 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/323/5916/934#otherarticles>

This article appears in the following **subject collections**:

Pharmacology, Toxicology

http://www.sciencemag.org/cgi/collection/pharm_tox

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

implications for optimizing plant breeding and cultivation strategies.

References and Notes

1. F. Laibach, *Arabidopsis Information Service 015* (1965).
2. L. Thompson, *J. Ecol.* **82**, 63 (1994).
3. Y. Kobayashi, D. Weigel, *Genes Dev.* **21**, 2371 (2007).
4. I. Ausin, C. Alonso-Blanco, J. M. Martinez-Zapater, *Int. J. Dev. Biol.* **49**, 689 (2005).
5. I. Baurle, C. Dean, *Cell* **125**, 655 (2006).
6. M. Koornneef, C. J. Hanhart, J. H. van der Veen, *Mol. Gen. Genet.* **229**, 57 (1991).
7. J. Putterill, F. Robson, K. Lee, R. Simon, G. Coupland, *Cell* **80**, 847 (1995).
8. R. N. Wilson, J. W. Heckman, C. R. Somerville, *Plant Physiol.* **100**, 403 (1992).
9. M. A. Blazquez, J. H. Ahn, D. Weigel, *Nat. Genet.* **33**, 168 (2003).
10. S. M. Welch, J. L. Roe, Z. Dong, *Agron. J.* **95**, 71 (2003).
11. J. Lempe *et al.*, *PLoS Genet.* **1**, e6 (2005).
12. S. Balasubramanian, S. Sureshkumar, J. Lempe, D. Weigel, *PLoS Genet.* **2**, e106 (2006).
13. R. J. Schmitz, R. M. Amasino, *Biochim. Biophys. Acta* **1769**, 269 (2007).
14. S. Sung, R. M. Amasino, *Nature* **427**, 159 (2004).
15. E. S. Dennis, W. J. Peacock, *Curr. Opin. Plant Biol.* **10**, 520 (2007).
16. G. G. Simpson, C. Dean, *Science* **296**, 285 (2002).
17. S. D. Michaels, Y. H. He, K. C. Scortecchi, R. M. Amasino, *Proc. Natl. Acad. Sci. U.S.A.* **100**, 10102 (2003).
18. See supporting material on Science Online.
19. T. Mizoguchi *et al.*, *Plant Cell* **17**, 2255 (2005).
20. J. H. Jung *et al.*, *Plant Cell* **19**, 2736 (2007).
21. I. Lee, R. M. Amasino, *Plant Physiol.* **108**, 157 (1995).
22. S. D. Michaels, R. M. Amasino, *Plant Cell* **13**, 935 (2001).
23. L. T. Burghardt *et al.*, in *19th International Conference on Arabidopsis Research* (Montreal, 23–27 July 2008), abstract ICAR802.
24. G. S. McMaster *et al.*, *Ann. Bot.* **102**, 561 (2008).
25. S. M. Welch, Z. Dong, J. L. Roe, *Aust. J. Agric. Res.* **56**, 919 (2005).
26. J. P. De Melo-Abreu *et al.*, *Agric. For. Meteorol.* **125**, 117 (2004).
27. G. T. Howe *et al.*, *Can. J. Bot.* **81**, 1247 (2003).
28. H. Böhlenius *et al.*, *Science* **312**, 1040 (2006); published online 3 May 2006 (10.1126/science.1126038).
29. J. Cockram *et al.*, *J. Exp. Bot.* **58**, 1231 (2007).
30. We thank M. Blazquez, C. Dean, M. Hoffmann, M. Koornneef, H. Kuittinen, and O. Savolainen for hosting these experiments at the European field sites; L. Albertson, L. Burghardt, C. Cooper, M. F. Cooper, E. Josephs, A. Lockwood, S. Myllylä, C. Oakley, R. Palmer, S. Rudder, and E. vonWettberg for assistance with planting and censusing; B. Robertson, M. Gosling, C. Lister, F. Eikelmann, G. Leuffen, W. Schuchert, T. Kauppila, and H. Eissner for technical and field assistance; R. Griffin, J. Kreikemeier, and G. Ragusa for assistance with the chamber experiment; A. Heim for contributions to the processing of temperature data; R. Amasino and J. Martinez-Zapater for sharing lines; S. Ruzsa of the Puruggan lab, who bulked lines for the experiments; and R. Amasino and C. Dean for manuscript comments. Supported by NSF Frontiers in Integrative Biological Research program grant EF-0425759 and an Alexander von Humboldt Research Award.

Supporting Online Material

www.sciencemag.org/cgi/content/full/1165826/DC1
Materials and Methods
Figs. S1 to S11
Tables S1 to S6
References

11 September 2008; accepted 19 November 2008
Published online 15 January 2009;
10.1126/science.1165826
Include this information when citing this paper.

The Hallucinogen *N,N*-Dimethyltryptamine (DMT) Is an Endogenous Sigma-1 Receptor Regulator

Dominique Fontanilla,¹ Molly Johannessen,² Abdol R. Hajjipour,³ Nicholas V. Cozzi,¹ Meyer B. Jackson,² Arnold E. Ruoho^{1*}

The sigma-1 receptor is widely distributed in the central nervous system and periphery. Originally mischaracterized as an opioid receptor, the sigma-1 receptor binds a vast number of synthetic compounds but does not bind opioid peptides; it is currently considered an orphan receptor. The sigma-1 receptor pharmacophore includes an alkylamine core, also found in the endogenous compound *N,N*-dimethyltryptamine (DMT). DMT acts as a hallucinogen, but its receptor target has been unclear. DMT bound to sigma-1 receptors and inhibited voltage-gated sodium ion (Na⁺) channels in both native cardiac myocytes and heterologous cells that express sigma-1 receptors. DMT induced hypermobility in wild-type mice but not in sigma-1 receptor knockout mice. These biochemical, physiological, and behavioral experiments indicate that DMT is an endogenous agonist for the sigma-1 receptor.

The sigma-1 receptor binds a broad range of synthetic compounds (1). It has long been suspected that the sigma-1 receptor is targeted by endogenous ligands, and several candidates have been proposed (2, 3). Although progesterone and other neuroactive steroids are known to bind sigma-1 receptors and regulate some of their functions (1, 4), they do not exhibit agonist properties on sigma-1-regulated ion channels in electrophysiological experiments (5).

Our search for a sigma receptor endogenous ligand (or ligands) was based on a variant of the

canonical sigma-1 receptor ligand pharmacophore (6), but with a more basic structure (Fig. 1A). Otherwise dissimilar sigma-1 receptor ligands possess a common *N*-substituted pharmacophore (Fig. 1A): an *N,N*-dialkyl or *N*-alkyl-*N*-aralkyl product, most easily recognized in the high-affinity sigma-1 receptor ligand, fenpropimorph (7). Similar chemical backbones can be derived from other sigma-1 receptor ligands such as haloperidol and cocaine (Fig. 1A). *N*-substituted trace amines harbor this sigma-1 receptor ligand pharmacophore, but their interactions with sigma receptors have not been determined. Of particular interest is the only known endogenous mammalian *N,N*-dimethylated trace amine, *N,N*-dimethyltryptamine (DMT) (8–10). In addition to being one of the active compounds in psychoactive snuffs (*yopo*, *ependá*) and sacramental teas (*ayahuasca*, *yagé*) used in native shamanic rituals in South America, DMT can be produced by enzymes in mammalian lung (11) and in rodent

brain (12). DMT has been found in human urine, blood, and cerebrospinal fluid (9, 13). Although there are no conclusive quantitative studies measuring the abundance of endogenous DMT because of its rapid metabolism (14), DMT concentrations can be localized and elevated in certain instances. Evidence suggests that DMT can be locally sequestered into brain neurotransmitter storage vesicles and that DMT production increases in rodent brain under environmental stress (8). Although a family of heterotrimeric GTP-binding protein (G protein)-coupled receptors (GPCRs) known as the trace amine receptors (TARs) was discovered in 2001 (15), only two members of this family respond to trace amines and have been renamed trace amine-associated receptors (TAARs) (16). Because other binding targets for trace amines and DMT are likely (8), we first examined the sigma-1 receptor binding affinities of the trace amines and their *N*-methylated and *N,N*-dimethylated counterparts.

Competition assays against the sigma-1 receptor-specific ligand, (+)-[³H]-pentazocine (10 nM), determined that the nonmethylated trace amines tryptamine, phenethylamine, and tyramine bound the sigma-1 receptor poorly (Fig. 1C), with dissociation constant (K_d) values of 431, 97.4, and >30,000 μ M, respectively. By contrast, the *N*-methylated and *N,N*-dimethylated derivatives of these compounds bound sigma-1 receptors more tightly, with a clear increase in affinity as the ligands approached the sigma-1 receptor ligand and pharmacophore (Fig. 1, A and B). With the exception of the *N*-methylated tyramines, this trend did not apply to the sigma-2 receptor, which differs pharmacologically and functionally from the sigma-1 receptor (Fig. 1C). Tryptamine, phenethylamine, and *N*-methyltryptamine had the highest sigma-2 receptor affinities, with K_d values of 4.91, 7.31, and 6.61 μ M, respectively. In contrast to sigma-1 receptors, *N*-methylation and *N,N*-

¹Department of Pharmacology, University of Wisconsin School of Medicine and Public Health, Madison, WI 53706, USA.

²Department of Physiology, University of Wisconsin School of Medicine and Public Health, Madison, WI 53706, USA.

³Pharmaceutical Research Laboratory, Department of Chemistry, Isfahan University of Technology, Isfahan 84156, IR Iran.

*To whom correspondence should be addressed. E-mail: aeruoho@wisc.edu

dimethylation of tryptamine and phenethylamine decreased sigma-2 receptor affinity (Fig. 1C).

We tested the ability of tryptamine, *N*-methyltryptamine, and DMT to block sigma receptor photolabeling in rat liver homogenates by two radioactive photoaffinity labels, the sigma-1 receptor–

specific cocaine derivative 3-¹²⁵Ijodo-4-azidococaine ([¹²⁵I]-IACoc) (17) (Fig. 2A) and the sigma-1 and sigma-2 receptor fenpropimorph derivative 1-*N*-(2,6'-dimethyl-morpholino)-3-(4-azido-3-[¹²⁵I]iodo-phenyl) propane ([¹²⁵I]IAF) (18) (Fig. 2B). Both of these compounds have been used to identify the drug

binding region of the sigma-1 receptor (18, 19). As anticipated, [¹²⁵I]-IACoc [sigma-1 *K*_d = 0.126 nM (17)] photolabeling of the 26-kD sigma-1 receptor (Fig. 2A) was protected best by DMT, with 61% protection by 50 μM DMT and almost 100% protection by 100 μM DMT. By contrast, tryptamine

Fig. 1. Sigma-1 receptor ligand pharmacophore and binding affinities. **(A)** A basic sigma-1 receptor ligand pharmacophore variant of Glennon *et al.* (6) was derived by removal of the red bonds from the sigma-1 receptor ligands fenpropimorph, haloperidol, and cocaine. **(B)** Competitive binding curves of tryptamine, *N*-methyltryptamine, and DMT, against the radioactive sigma-1 receptor ligand [³H]-(+)-pentazocine. Curves are shown as percent specific binding (5 μM haloperidol). **(C)** Sigma-1 and sigma-2 receptor *K*_d values of trace amines and their *N*-methylated and *N,N*-dimethylated derivatives (scheme S2). Included are SEM values (*n* = 3 binding experiments) and *R*² values for a nonlinear regression curve fit. Solid arrows denote the direction of increasing affinity.






Fig. 3. Sodium channel inhibition by DMT.


(A) In the presence or absence of 10 μM haloperidol, wild type (WT) or sigma-1 receptor knock-out (KO) mouse liver homogenates (200 $\mu\text{g}/\text{lane}$) were photolabeled with 1 nM [^{125}I]IAF. (B) Examples of I_{Na} evoked by steps from -80 to -10 mV in HEK293 or COS-7 cells expressing hNav1.5 channel in the absence (control, black), presence (DMT, red), and after wash out (recovery, blue) of 100 μM DMT. Average inhibition by DMT was determined by measuring peak I_{Na} . Bars represent mean \pm SEM ($n = 3$ cells). I_{Na} inhibition in HEK293 cells differed significantly from that in COS-7 cells ($*P < 0.03$). (C) Examples of I_{Na} evoked as described in (B) in neonatal cardiac myocytes from WT and KO mice in the absence (control, black), presence (DMT, red), and after wash out (recovery, blue) of 100 μM DMT. Current inhibition in WT was significantly different from that in KO ($*P < 0.002$, $n = 7$ neonatal cardiac myocytes).



and *N*-methyltryptamine protected minimally against sigma-1 receptor [^{125}I]-IACoc photolabeling, even at these high concentrations (Fig. 2A). Similarly, [^{125}I]IAF photolabeling of the sigma-1 [$K_d = 194$ nM (18)] receptor showed that DMT was the most potent protector. Ten micromolar DMT provided 31% protection, whereas 50 and 100 μM DMT provided 43 and 69% protection, respectively (Fig. 2B). With the exception of *N*-methyltryptamine, protection of [^{125}I]IAF sigma-2 [$K_d = 2780$ nM (18)] receptor photolabeling paralleled the sigma-2 binding data. Tryptamine afforded the greatest protection of sigma-2 receptor photolabeling, with values of 47, 78, and 79% for 10, 50, and 100 μM , respectively (Fig. 2B).

An important biological activity of sigma receptor activation is the inhibition of ion channels, which operates through protein-protein interactions without mediation by G proteins and protein kinases (20–22). In addition to modulating various types of voltage-activated K^+ channels (21, 23, 24), the sigma-1 receptor associates with the $\text{Kv}1.4$ K^+ channel in posterior pituitary nerve terminals, as well as in *Xenopus* oocytes (22). Sigma receptor ligands also modulate N-, L-, P/Q-, and R-type Ca^{2+} channels in rat sympathetic and parasympathetic neurons (25). Sigma receptor ligands modulate cardiac voltage-gated Na^+ channels (hNav1.5) in human embryonic kidney 293 (HEK293) cells, COS-7 cells, and neonatal mouse cardiac myocytes (26). To evaluate the capacity of DMT to induce physiological responses by binding to sigma receptors, we examined the action of DMT on voltage-activated Na^+ current. Patch-clamp recordings from HEK293 cells stably expressing the human cardiac Na^+ channel hNav1.5 revealed voltage-activated Na^+ currents (I_{Na}) in response to voltage steps from -80 to -10 mV (Fig. 3B). Application of 100 μM DMT inhibited I_{Na} by $62 \pm 3\%$ ($n = 3$ HEK293 cells), which reversed upon DMT removal. With hNav1.5 transiently transfected into COS-7 cells,

Fig. 4. DMT-induced hypermobility abrogated in the sigma-1 KO mouse. (A) Distances traveled by WT and KO mice were measured in an open-field assay in 5-min increments. Pargyline was injected 2 hours before DMT or vehicle (Veh) ip injection. Bars represent mean \pm SEM ($n = 8$ to 14 mice). WT mice showed a significant ($***P < 0.0001$) increase in mobility in response to DMT as compared to KO mice. (B) Total distance traveled over 30 min after DMT, vehicle (Veh), or methamphetamine (Meth, $n = 6$ mice) injection in WT and KO mice. (C) Methamphetamine serves as a positive control for hypermobility in KO mice.



100 μM DMT inhibited I_{Na} by only $22 \pm 4\%$ ($n = 3$ COS-7 cells), but photolabeling has shown that these cells have much lower concentrations of endogenous sigma-1 receptors compared to HEK293 cells (fig. S1 and Fig. 3B). The difference between DMT inhibition of I_{Na} in HEK293 and COS-7

cells (Fig. 3B, $P < 0.03$) thus demonstrates the dependence of I_{Na} inhibition on sigma-1 receptors. Experiments in cardiac myocytes demonstrated the same DMT action in a native preparation (Fig. 3C) and enabled further demonstration of sigma-1 receptor dependence by using a sigma-1 receptor

knockout mouse (27). [¹²⁵I]IAF photolabeling of liver homogenates from wild-type (WT) and sigma-1 receptor knockout (KO) mice indeed showed the absence of sigma-1 receptor (26 kD) in the KO samples (Fig. 3A). In WT neonatal cardiac myocytes, 100 μM DMT reversibly inhibited I_{Na} by 29 ± 3% ($n = 7$ WT myocytes), whereas I_{Na} was reduced by only 7 ± 2% ($n = 7$ KO myocytes) in KO myocytes (Fig. 3C, $P < 0.002$).

Both DMT and sigma receptor ligands influence animal behavior. DMT injection induces hypermobility in rodents concurrently treated with the monoamine oxidase inhibitor pargyline (28), and this action is not antagonized by blockers of dopamine or serotonin receptors, but is potentially inhibited by haloperidol (28). Although haloperidol is thought to act in part through the dopamine D₂ receptor system, it is also a potent sigma-1 receptor agonist [sigma-1 inhibition constant (K_i) = 3 nM (29); sigma-2 K_i = 54 nM (29)] when inhibiting voltage-gated ion channels (5, 25). Haloperidol reduces brain concentrations of DMT (8) and DMT inhibits haloperidol binding in brain tissue more robustly than the dopamine agonist apomorphine (8). On the basis of these findings, which were discovered before sigma receptor identification, DMT has been hypothesized to act through an unknown “hallucinogen” receptor (8). We confirmed results (28) that intraperitoneal (ip) administration of DMT (2 mg per kilogram of body weight) 2 hours after pargyline (75 mg/kg, ip) injection induced hypermobility in WT mice (7025 ± 524.1 cm, $n = 12$ WT mice) in an open-field assay. Identical drug treatments in sigma-1 receptor KO mice had no hypermobility action (2328 ± 322.9 cm, $n = 12$ KO mice, $P < 0.0001$; Fig. 4, A and B). This result is particularly important to our understanding of sigma-1 receptor biological function because the KO mice are viable and fertile (27). The sigma-1 receptor dependence of DMT-induced hypermobility parallels that induced by the sigma-1 receptor ligand (+)-SKF10047 in WT but not in KO mice (27). As a positive control, methamphetamine, which is thought to act through catecholaminergic systems, induced hypermobility in both WT and KO mice (3 mg/kg, ip, $n = 6$ mice; Fig. 4, B and C) with a reduced onset rate compared with that seen for DMT (Fig. 4, A and C). This indicates that behavioral actions of DMT depend on the sigma-1 receptor, which may provide an alternative research area for psychiatric disorders that have not been linked to dopamine or *N*-methyl-D-aspartate systems.

The binding, biochemical, physiological, and behavioral studies reported here all support the hypothesis that DMT acts as a ligand for the sigma-1 receptor. On the basis of our binding results and the sigma-1 receptor pharmacophore, endogenous trace amines and their *N*-methyl and *N,N*-dimethyl derivatives are likely to serve as endogenous sigma receptor regulators. Moreover, DMT, the only known mammalian *N,N*-dimethylated trace amine, can activate the sigma-1 receptor to modulate Na⁺ channels. The recent discovery that the sigma-1 receptor functions as a molecular chaperone (30) may be

relevant, because sigma-1 receptors, which are observed in the endoplasmic reticulum, associate with plasma membrane Kv 1.4 channels (22) and may serve as a molecular chaperone for ion channels. Furthermore, the behavioral effect of DMT may be due to activation or inhibition of sigma-1 receptor chaperone activity instead of, or in addition to, DMT/sigma-1 receptor modulation of ion channels. These studies thus suggest that this natural hallucinogen could exert its action by binding to sigma-1 receptors, which are abundant in the brain (1, 27). This discovery may also extend to *N,N*-dimethylated neurotransmitters such as the psychoactive serotonin derivative *N,N*-dimethylserotonin (bufotenine), which has been found at elevated concentrations in the urine of schizophrenic patients (10). The finding that DMT and sigma-1 receptors act as a ligand-receptor pair provides a long-awaited connection that will enable researchers to elucidate the biological functions of both of these molecules.

References and Notes

1. T. Hayashi, T. P. Su, *CNS Drugs* **18**, 269 (2004).
2. P. Bouchard *et al.*, *Eur. J. Neurosci.* **7**, 1952 (1995).
3. T. P. Su, A. D. Weissman, S. Y. Yeh, *Life Sci.* **38**, 2199 (1986).
4. T. P. Su, E. D. London, J. H. Jaffe, *Science* **240**, 219 (1988).
5. R. A. Wilke *et al.*, *J. Physiol.* **517**, 391 (1999).
6. R. A. Glennon *et al.*, *J. Med. Chem.* **37**, 1214 (1994).
7. F. F. Moebius, R. J. Reiter, M. Hanner, H. Glossmann, *Br. J. Pharmacol.* **121**, 1 (1997).
8. S. A. Barker, J. A. Monti, S. T. Christian, *Int. Rev. Neurobiol.* **22**, 83 (1981).
9. F. Franzen, H. Gross, *Nature* **206**, 1052 (1965).
10. M. S. Jacob, D. E. Presti, *Med. Hypotheses* **64**, 930 (2005).
11. J. Axelrod, *Science* **134**, 343 (1961).
12. J. M. Saavedra, J. Axelrod, *Science* **175**, 1365 (1972).
13. J. M. Beaton, P. E. Morris, *Mech. Ageing Dev.* **25**, 343 (1984).
14. S. A. Burchett, T. P. Hicks, *Prog. Neurobiol.* **79**, 223 (2006).
15. B. Borowsky *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **98**, 8966 (2001).
16. L. Lindemann *et al.*, *Genomics* **85**, 372 (2005).
17. J. R. Kahoun, A. E. Ruoho, *Proc. Natl. Acad. Sci. U.S.A.* **89**, 1393 (1992).
18. A. Pal *et al.*, *Mol. Pharmacol.* **72**, 921 (2007).
19. Y. Chen, A. R. Hajipour, M. K. Sievert, M. Arbabian, A. E. Ruoho, *Biochemistry* **46**, 3532 (2007).
20. P. J. Lupardus *et al.*, *J. Physiol.* **526**, 527 (2000).
21. H. Zhang, J. Cuevas, *J. Pharmacol. Exp. Ther.* **313**, 1387 (2005).
22. E. Aydar, C. P. Palmer, V. A. Klyachko, M. B. Jackson, *Neuron* **34**, 399 (2002).
23. R. A. Wilke *et al.*, *J. Biol. Chem.* **274**, 18387 (1999).
24. C. Kennedy, G. Henderson, *Neuroscience* **35**, 725 (1990).
25. H. Zhang, J. Cuevas, *J. Neurophysiol.* **87**, 2867 (2002).
26. M. A. Johannesssen, A. Ramos-Serrano, S. Ramachandran, A. E. Ruoho, M. B. Jackson, “Sigma receptor modulation of voltage-dependent sodium channels,” Program No. 466.22, Annual Neuroscience Meeting, San Diego, CA, 5 November 2007.
27. F. Langa *et al.*, *Eur. J. Neurosci.* **18**, 2188 (2003).
28. P. Jenner, C. D. Marsden, C. M. Thanki, *Br. J. Pharmacol.* **69**, 69 (1980).
29. R. R. Matsumoto, B. Pouw, *Eur. J. Pharmacol.* **401**, 155 (2000).
30. T. Hayashi, T. P. Su, *Cell* **131**, 596 (2007).
31. We thank the Corinna Burger laboratory for use of their mouse behavior equipment, and A. Paul and T. Mavlyutov for providing [¹²⁵I]IAF and [¹²⁵I]IACoc, respectively. Supported by the Molecular and Cellular Pharmacology (MCP) Graduate Program training grant from NIH T32 GM08688 and by the NIH Ruth L. Kirschstein National Research Service Award (NRSA) (F31 DA022932) from the National Institute on Drug Abuse (to D.F.). This work was funded by NIH grants R01 MH065503 (to A.E.R.) and NS30016 (to M.B.J.).

Supporting Online Material

www.sciencemag.org/cgi/content/full/323/5916/934/DC1
Materials and Methods
Fig. S1 and scheme S2
References

18 September 2008; accepted 10 December 2008
10.1126/science.1166127

When Your Gain Is My Pain and Your Pain Is My Gain: Neural Correlates of Envy and Schadenfreude

Hidehiko Takahashi,^{1,2,3*} Motoichiro Kato,⁴ Masato Matsuura,² Dean Mobbs,⁵ Tetsuya Suhara,¹ Yoshiro Okubo⁶

We often evaluate the self and others from social comparisons. We feel envy when the target person has superior and self-relevant characteristics. Schadenfreude occurs when envied persons fall from grace. To elucidate the neurocognitive mechanisms of envy and schadenfreude, we conducted two functional magnetic resonance imaging studies. In study one, the participants read information concerning target persons characterized by levels of possession and self-relevance of comparison domains. When the target person's possession was superior and self-relevant, stronger envy and stronger anterior cingulate cortex (ACC) activation were induced. In study two, stronger schadenfreude and stronger striatum activation were induced when misfortunes happened to envied persons. ACC activation in study one predicted ventral striatum activation in study two. Our findings document mechanisms of painful emotion, envy, and a rewarding reaction, schadenfreude.

Envy is one of the seven biblical sins, the Shakespearean “green-eyed monster,” and what Bertrand Russell (1) called an unfortunate facet of human nature. It is an irrational, unpleasant feeling and a “painful emotion” (2)

characterized by feelings of inferiority and resentment produced by an awareness of another's superior quality, achievement, or possessions (3). Understanding envy is important because of its broad implications, ranging from individual mat-